



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

II. Solution by G. B. M. ZERR, A. M., Ph. D., 4243 Girard Avenue, Philadelphia, Pa.

General expressions for the sides of duplicate right-triangles having the same hypotenuse are

$$x = (p^2 - q^2)(r^2 - s^2) + 4pqrs, \quad y = 2rs(p^2 - q^2) - 2pq(r^2 - s^2),$$

$$z = (r^2 - s^2)(p^2 + q^2), \quad w = 2rs(p^2 + q^2).$$

$$\therefore x^2 + y^2 = z^2 + w^2 = [(r^2 + s^2)(p^2 + q^2)]^2, \quad \text{Let } p=2, q=1.$$

$$\therefore x = 3(r^2 - s^2) + 8rs, \quad y = 6rs - 4(r^2 - s^2), \quad z = 5(r^2 - s^2), \quad w = 10rs.$$

$$x^2 + y^2 = z^2 + w^2 = [5(r^2 + s^2)]^2, \quad x^2 - w^2 = z^2 - y^2 = 9r^4 + 48r^3s - 54r^2s^2 - 48rs^3 + 9s^4 = \square. \quad \text{This is a square when } r = \frac{1}{2}s.$$

$$\therefore x = \frac{689s^2}{64}, \quad y = \frac{161s^2}{36}, \quad z = \frac{725s^2}{144}, \quad w = \frac{85s^2}{6}.$$

$$\therefore x = 2067s^2, \quad y = 644s^2, \quad z = 725s^2, \quad w = 2040s^2.$$

$$x^2 + y^2 = z^2 + w^2 = (2165s^2)^2 = a^2, \quad x^2 - w^2 = z^2 - y^2 = (333s^2)^2 = b^2.$$

A solution of this problem is given in J. D. Williams' *Algebra*, page 419. He starts with $a^2 = b^2 + f^2 = c^2 + e^2$, and $b^2 - c^2 = d^2 = e^2 - f^2$. Then he is to make $a^2 - b^2$ a square, $a^2 - c^2$ a square, and $b^2 - f^2$ a square. He assumes $a^2 = (p^2 + q^2)(r^2 + s^2)$, $b = pr \pm qs$, $c = ps \pm qr$. Then he assumes $r = pm - qn$, $s = pn + qm$. He finally arrives at the conclusion that $a = 697$, $b = 680$, $f = 153$, $c = 672$, $e = 185$, a set of erroneous values, as Dr. Zerr has pointed out. It is likely that Williams' solution may be carried out so that a set of correct values may be obtained. Williams proposed this problem in 1832 as a challenge problem to the mathematicians of the United States. ED. F.

144. Proposed by JOHN D. WILLIAMS (being the ninth of his 14 challenge problems proposed in 1832).

Make $(m^2 + n^2)^2 x^2 \pm (m^2 + n^2)x = \square$, $(m^2 - n^2)^2 x^2 \pm (m^2 - n^2)x = \square$, and $4m^2 n^2 x^2 \pm 2mnx = \square$.

Solution by G. B. M. ZERR, A. M., Ph. D., 4243 Girard Avenue, Philadelphia, Pa.

$$\text{Let } m^2 + n^2 = p, \quad m^2 - n^2 = q, \quad 2mn = r.$$

$$\therefore p^2 x^2 \pm px = \square, \quad q^2 x^2 \pm qx = \square, \quad r^2 x^2 \pm rx = \square \dots (1, 2, 3).$$

$$\text{Let } p^2 x^2 \pm px = a^2 x^2; \therefore x = p/(a^2 - p^2).$$

This value of x in (2) and (3) gives

$$q^2 p^2 + qp(a^2 - p^2) = \square, \quad r^2 p^2 + rp(a^2 - p^2) = \square \dots (4, 5).$$

$$\text{Let } q^2 p^2 + qp(a^2 - p^2) = [pq - b(a - p)]^2.$$

$$\therefore (a - p) = \frac{2pq(b + p)}{b^2 - pq}. \quad \text{This value of } a \text{ in (5) gives}$$

$$r^2 b^4 + 4b^3 pqr + 2b^2 pqr(2p + 2q - r) + 4bp^2 q^2 r + p^2 q^2 r^2 = \square \\ = (rb + 2bpq - pqr)^2, \quad \text{suppose.}$$

$$\therefore b = \frac{2pqr}{pr + qr - pq}; \quad x = \frac{p}{a^2 - p^2} = \frac{(b^2 - pq)^2}{4bpq(b + p)(b + q)}.$$

$$\therefore x = \frac{-(pr + qr - pq)^2 - 4pqr^2}{8pqr(pr + qr - pq)(pq - pr + qr)(pq + pr - qr)}.$$

$$\therefore x = \frac{-[(4m^3n - m^4 + n^4)^2 - 16m^2n^2(m^4 - n^4)]^2}{16mn(m^4 - n^4)(4m^3n - m^4 + n^4)(m^4 - n^4 - 4mn^3)(m^4 - n^4 + 4mn^3)}$$

$$= \pm \frac{A^2}{16mn(m^4 - n^4)B}, \text{ suppose.}$$

x is \pm according as px , qx , rx is \mp .

$$\therefore (m^2 + n^2)x^2 \pm (m^2 + n^2)x$$

$$= \left[\frac{A}{16mn(m^2 - n^2)B} \right]^2 [16m^2n^4(n^2 - 2m^2) + (8mn^3 + n^4 - m^4)(m^4 - n^4)]^2. (6).$$

$$(m^2 - n^2)x^2 \pm (m^2 - n^2)x$$

$$= \left[\frac{A}{16mn(m^2 + n^2)B} \right]^2 [16m^2n^4(n^2 + 2m^2) - (8mn^3 + m^4 - n^4)(m^4 - n^4)]^2. (7).$$

$$4m^2n^2x^2 \pm 2mnx$$

$$= \left[\frac{A}{8(m^4 - n^4)B} \right]^2 [3(m^4 - n^4)^2 - 8m^3n(m^4 - n^4) - 16m^2n^6]^2. (8).$$

m and n can have any values that make B positive. Let $m=2$, $n=1$;
 $A=671$, $B=2737$.

$$(6) \text{ gives } (m^2 + n^2)^2 x^2 \pm (m^2 + n^2)x = (290543/262752)^2.$$

$$(7) \text{ gives } (m^2 - n^2)^2 x^2 \pm (m^2 - n^2)x = (74481/437920)^2.$$

$$(8) \text{ gives } 4m^2n^2x^2 \pm 2mnx = (234179/328440)^2.$$

AVERAGE AND PROBABILITY.

191. Proposed by J. EDWARD SANDERS, Reinersville, Ohio.

Two random lines cut a given circle. What is the chance that they intersect within the circle?

II. Solution by G. B. M. ZERR, A. M., Ph. D., 4243 Girard Avenue, Philadelphia, Pa.

Let AB , CD be the random lines, $\angle AOH = \theta$,
 $\angle COE = \phi$, $\angle EOH = \psi$.

The limits of θ are 0 and $\frac{1}{2}\pi$; of ϕ , 0 and θ ; of ψ , $\theta - \phi$ and $\theta + \phi$ for favorable cases, and 0 and π for total cases.

Hence the chance is

